

SPLITTING OFF RATIONAL PARTS IN HOMOTOPY TYPES

NORIO IWASE[†] AND NOBUYUKI ODA^{††}

ABSTRACT. It is known algebraically that any abelian group is a direct sum of a divisible group and a reduced group (See Theorem 21.3 of [5]). In this paper, conditions to split off rational parts in homotopy types from a given space are studied in terms of a variant of Hurewicz map, say $\bar{p}: [S_{\mathbb{Q}}^n, X] \rightarrow H_n(X; \mathbb{Z})$ and generalized Gottlieb groups. This yields decomposition theorems on rational homotopy types of Hopf spaces, T -spaces and Gottlieb spaces, which has been known in various situations, especially for spaces with finiteness conditions.

INTRODUCTION

The Gottlieb group is introduced by Gottlieb [6, 7] and the generalized Gottlieb set is introduced by Varadarajan [17]. Dula and Gottlieb obtained a general result on splitting a Hopf space off from a fibration as Theorem 1.3 of [4].

In this paper, we work in the category of spaces having homotopy types of CW complexes with base points and pointed continuous maps. A relation $f \sim g$ indicates a pointed homotopy relation of maps f and g and a relation $X \simeq Y$ indicates a homotopy equivalence relation of spaces X and Y . We also denote by $[X, Y]$ the set of pointed homotopy classes of maps from X to Y .

We adopt some more conventional notations: $X_{\mathbb{Q}}$ stands for the rationalization of a space X , $K(\pi, n)$ for the Eilenberg-Mac Lane space of type (π, n) , $G(V, X)$ for the generalized Gottlieb subset of $[V, X]$ and $H_n(X)$ for $H_n(X; \mathbb{Z})$. We introduce a variant of Hurewicz map $\bar{p}: [S_{\mathbb{Q}}^n, X] \rightarrow H_n(X)$ by $\bar{p}(\alpha) = \alpha_*([S^n] \otimes 1)$ for $\alpha \in [S_{\mathbb{Q}}^n, X]$, where α_* is the homomorphism given by $\alpha_*: H_n(S^n) \otimes \mathbb{Q} = H_n(S_{\mathbb{Q}}^n) \rightarrow H_n(X)$. Our main result is described as follows:

Theorem 2.2. *Let $R = \bigoplus_{\lambda \in \Lambda} \mathbb{Q}$ be a $\mathbb{Q}$ -vector space of dimension $\#\Lambda \leq \infty$. Let X be 0-connected and $R \subset \bar{p}(G(S_{\mathbb{Q}}^n, X)) \subseteq H_n(X)$, $n \geq 2$. Then X decomposes as*

$$X \simeq Y \times K(R, n).$$

Theorem 2.2 gives unified proof to the splitting phenomena on rational G-space, T -space and Hopf space without assuming any finiteness conditions, which are proved under various situations by a number of authors: Scheerer [16] obtained decomposition theorems of rational Hopf spaces without assuming the finite type assumptions. Oprea [15] obtained decomposition theorems by using minimal model method in rational homotopy theory. Aguadé [1] obtained a decomposition theorems on rational T -spaces of finite type.

Date: February 1, 2008.

[†]supported by the Grant-in-Aids for Scientific Research #14654016 from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

^{††}supported by the Grant-in-Aids for Scientific Research #15340025 from the Japan Society for the Promotion of Science.

1. PRELIMINARIES

We regard the one point union $X \vee Y$ of spaces X and Y as a subspace $X \times * \cup * \times Y$ of the product space $X \times Y$ with the inclusion map $j : X \vee Y \rightarrow X \times Y$. For any collection of a finitely or infinitely many spaces X_λ ($\lambda \in \Lambda$), we denote the *wedge sum* (or one point union) by $\bigvee_{\lambda \in \Lambda} X_\lambda$ and the *direct sum* (or weak product) by $\bigoplus_{\lambda \in \Lambda} X_\lambda = \{(x_\lambda) \in \prod_{\lambda \in \Lambda} X_\lambda \mid x_\lambda = * \text{ except for finitely many } \lambda\}$. Then we have $\bigvee_{\lambda \in \Lambda} X_\lambda \subset \bigoplus_{\lambda \in \Lambda} X_\lambda$, where $\bigoplus_{\lambda \in \Lambda} X_\lambda$ is a dense subset of the product space $\prod_{\lambda \in \Lambda} X_\lambda$ and has the weak topology with respect to finite products of X_λ 's.

We can apply rationalization or $\mathbb{Q}$ -localization to any 0-connected nilpotent spaces (cf. [3], [9] or [13]). The rationalization $\ell_{\mathbb{Q}} : X \rightarrow X_{\mathbb{Q}}$, or simply $X_{\mathbb{Q}}$ does exist for such spaces X such that $\ell_{\mathbb{Q}}$ induces the following isomorphisms:

$$\pi_n(X_{\mathbb{Q}}) \cong \pi_n(X) \otimes \mathbb{Q} \quad \text{and} \quad H_n(X_{\mathbb{Q}}) \cong H_n(X) \otimes \mathbb{Q}$$

for any integer $n \geq 1$. Moreover the universality of rationalization yields a bijection

$$\ell_{\mathbb{Q}}^* : [X_{\mathbb{Q}}, Y_{\mathbb{Q}}] \cong [X, Y_{\mathbb{Q}}]$$

for any such spaces X and Y . The rationalization enjoys the following fact.

Fact 1.1. (1) $S_{\mathbb{Q}}^{2m+1} \simeq K(\mathbb{Q}, 2m+1)$ for any integer $m \geq 0$.
 (2) $\Omega(S_{\mathbb{Q}}^{2m+1}) \simeq (\Omega S^{2m+1})_{\mathbb{Q}} \simeq K(\mathbb{Q}, 2m)$ for any integer $m \geq 1$.
 (3) $(X_{\infty})_{\mathbb{Q}} \simeq (X_{\mathbb{Q}})_{\infty}$ for X a 0-connected nilpotent space of finite type.

Proof. (1) and (2) are well-known. We give here a brief explanation for (3): The suspension functor Σ and the loop functor Ω enjoys the properties $\Sigma(X_{\mathbb{Q}}) \simeq (\Sigma X)_{\mathbb{Q}}$ for any 0-connected space X and $\Omega(X_{\mathbb{Q}}) \simeq (\Omega X)_{\mathbb{Q}}$ for any 1-connected space X . Let X_{∞} be the James reduced product space of a 0-connected space X of finite type, so that $X_{\infty} \simeq \Omega(\Sigma X)$ by James [10]. Then it follows that $(X_{\infty})_{\mathbb{Q}} \simeq (\Omega(\Sigma X))_{\mathbb{Q}} \simeq \Omega(\Sigma(X_{\mathbb{Q}})) \simeq (X_{\mathbb{Q}})_{\infty}$. $\square$

We state two propositions to be used in the proof of the main theorem.

Proposition 1.2. *Let X be a 0-connected space of finite type and $f : X \rightarrow Y$ a map. If $f \in G(X, Y)$, then there is an extension $\bar{f} : X_{\infty} \rightarrow Y$ of f such that $\bar{f} \in G(X_{\infty}, Y)$.*

Proof. We may assume that there is a map $\mu : Y \times X \rightarrow Y$ such that $\mu|_{Y \times \{*\}} = 1_Y : Y \rightarrow Y$ and $\mu|_{\{*\} \times X} = f : X \rightarrow Y$. We put $\mu_1 = \mu$ and, for any n we define

$$\mu_n = \mu \circ (\mu_{n-1} \times 1_X) : Y \times X^n = (Y \times X^{n-1}) \times X \rightarrow Y \times X \rightarrow Y$$

by induction on n . Then we observe that μ_n factors through $Y \times X^n \rightarrow Y \times X_n$. $\square$

Proposition 1.3. *Let $\alpha_\lambda : X_\lambda \rightarrow Z$ be a map for any $\lambda \in \Lambda$. If $\alpha_\lambda \in G(X_\lambda, Z)$ for each $\lambda \in \Lambda$, then the map $\alpha : \bigvee_{\lambda \in \Lambda} X_\lambda \rightarrow Z$ defined by $\alpha|_{X_\lambda} = \alpha_\lambda : X_\lambda \rightarrow Z$ can be extended to a map $\bar{\alpha} : \bigoplus_{\lambda \in \Lambda} X_\lambda \rightarrow Z$ with $\bar{\alpha} \in G(\bigoplus_{\lambda \in \Lambda} X_\lambda, Z)$.*

Proof. Since each X_λ has a homotopy type of a CW complex, we may assume that there is a map $\mu_\lambda : Z \times X_\lambda \rightarrow Z$ such that $\mu_\lambda|_{\{*\} \times X_\lambda} = \alpha_\lambda : X_\lambda \rightarrow Z$ and $\mu_\lambda|_{Z \times \{*\}} = 1_Z : Z \rightarrow Z$ for each $\lambda \in \Lambda$. For any n and $\lambda_1, \lambda_2, \dots, \lambda_n$, we define

$$\mu_{\lambda_1, \dots, \lambda_n} = \mu_{\lambda_n} \circ (\mu_{\lambda_1, \dots, \lambda_{n-1}} \times 1_{X_{\lambda_n}}) : Z \times (X_{\lambda_1} \times \dots \times X_{\lambda_{n-1}} \times X_{\lambda_n}) \rightarrow Z$$

by induction on n . For any index set Λ , we assume that Λ is totally-ordered. Then we can easily observe that the collection of maps $\mu_{\lambda_1, \dots, \lambda_n}$ defines a pairing $\mu : Z \times (\bigoplus_{\lambda \in \Lambda} X_\lambda) \rightarrow Z$ with axes $(1_Z, \bar{\alpha})$ (cf. [14]). $\square$

2. PROOF OF THE MAIN RESULT

Proposition 2.1. *Let P be an idempotent endomorphism of $H_n(X)$, $n \geq 2$. Suppose that $R = \text{im } P \subseteq H_n(X)$ is a rational vector space and is in $\text{im } \bar{\rho}$. Then we have maps $\alpha : S^n(R) \rightarrow X$ and $\beta : X \rightarrow K(R, n)$ such that*

$$\beta \circ \alpha \sim \iota_R^n : S^n(R) \rightarrow K(R, n), \text{ and}$$

$$P = \alpha_* \circ (\iota_{R*}^n)^{-1} \circ \beta_* : H_n(X) \rightarrow H_n(K(R, n)) \xrightarrow{\cong} H_n(S^n(R)) \rightarrow H_n(X),$$

where $S^n(R)$ denotes the Moore space of type (R, n) and ι_R^n corresponds to the identity element in $\text{Hom}(R, R) = \text{Hom}(\pi_n(S^n(R)), \pi_n(K(R, n))) \cong [S^n(R), K(R, n)]$.

Proof. Let $\{\bar{\rho}(\alpha_\lambda) \mid \lambda \in \Lambda\}$ be a basis of $R = \text{im } P$, and hence $R \cong \bigoplus_{\lambda \in \Lambda} \mathbb{Q}$. Since $S^n(R) = \bigvee_{\lambda \in \Lambda} S_{\mathbb{Q}}^n$, we define $\alpha : S^n(R) \rightarrow X$ by its restrictions to all factors:

$$\alpha|_{S_{\mathbb{Q}}^n} = \alpha_\lambda : S_{\mathbb{Q}}^n \rightarrow X.$$

Since α_* is an isomorphism onto $R \subseteq H_n(X)$, we have its inverse $\phi : R \rightarrow H_n(S_{\mathbb{Q}}^n)$ so that $\phi \circ \alpha_* = \text{id}_{H_n(S_{\mathbb{Q}}^n)}$ and $\alpha_* \circ \phi = \text{id}_R$. Now we define a homomorphism $s : H_n(X) \rightarrow \text{im } P \cong H_n(S_{\mathbb{Q}}^n)$ by $s = \phi \circ P$: Since $\text{im } \alpha_*$ is in the image of an idempotent endomorphism P , we have $s \circ \alpha_* = \phi \circ P \circ \alpha_* = \phi \circ \alpha_* = \text{id}$. Also we have $\alpha_* \circ s = \alpha_* \circ \phi \circ P = P$. Thus s satisfies the following formulae:

$$s \circ \alpha_* = \text{id} : H_n(S^n(R)) \rightarrow H_n(S^n(R)),$$

$$\alpha_* \circ s = P : H_n(X) \rightarrow H_n(X).$$

Let us recall that α induces the following commutative diagram:

$$(2.1) \quad \begin{array}{ccc} [X, K(R, n)] & \xrightarrow[\cong]{\Psi'} & \text{Hom}(H_n(X), H_n(K(R, n))) \\ \alpha_* \downarrow & & \downarrow (\alpha_*)^* \\ [S^n(R), K(R, n)] & \xrightarrow[\cong]{\Psi} & \text{Hom}(H_n(S^n(R)), H_n(K(R, n))), \end{array}$$

where Ψ and Ψ' are homomorphisms defined by taking the n -th homology groups, and are isomorphisms by the universal coefficient theorem. Since Ψ' is an isomorphism, we define β to be the unique element $\Psi'^{-1}(\iota_{R*}^n \circ s)$ so that $\beta_* = \iota_{R*}^n \circ s$.

Firstly by $P = \alpha_* \circ s$, we have $P = \alpha_* \circ s = \alpha_* \circ (\iota_{\mathbb{Q}*}^n)^{-1} \circ \beta_*$.

Next we show $\beta \circ \alpha \sim \iota_{\mathbb{Q}}^n$. By the commutativity of the diagram (2.1), we have

$$\Psi(\alpha^*(\beta)) = (\alpha_*)^* \circ \Psi'(\beta) = (\alpha_*)^*(\iota_{\mathbb{Q}*}^n \circ s) = \iota_{\mathbb{Q}*}^n \circ s \circ \alpha_* = \iota_{\mathbb{Q}*}^n = \Psi(\iota_{\mathbb{Q}}^n).$$

Since Ψ is an isomorphism, we also have $\beta \circ \alpha = \alpha^*(\beta) \sim \iota_{\mathbb{Q}}^n$. $\square$

Let us recall that $G(S_{\mathbb{Q}}^n, X) \subset [S_{\mathbb{Q}}^n, X] \xrightarrow{\bar{\rho}} H_n(X)$. In the following theorem, we do *not* assume that X is rationalized nor that X is $(n-1)$ -connected.

Theorem 2.2. *Let $R = \bigoplus_{\lambda \in \Lambda} \mathbb{Q}$ be a $\mathbb{Q}$ -vector space of dimension $\#\Lambda \leq \infty$. Let X be 0-connected and $R \subset \bar{\rho}(G(S_{\mathbb{Q}}^n, X)) \subseteq H_n(X)$, $n \geq 2$. Then X decomposes as*

$$X \simeq Y \times K(R, n).$$

Proof. Since a divisible submodule R is a direct summand of $H_n(X)$, there is an idempotent endomorphism $P : H_n(X) \rightarrow H_n(X)$ with $\text{im } P = R$. We fix a basis of R as $\{\bar{\rho}(\alpha_\lambda) \mid \alpha_\lambda \in G(S_{\mathbb{Q}}^n, X), \lambda \in \Lambda\}$.

By Proposition 2.1, there are maps $\alpha : S^n(R) \rightarrow X$, $\beta : X \rightarrow K(R, n)$ such that

$$\beta \circ \alpha \sim \iota_R^n : S^n(R) \rightarrow K(R, n),$$

$$P = \alpha_* \circ (\iota_R^n)^{-1} \circ \beta_* : H_n(X) \rightarrow H_n(K(R, n)) \xrightarrow{\cong} H_n(S^n(R)) \rightarrow H_n(X).$$

Then we extend the map α onto $K(R, n) \supseteq S^n(R)$ as $\bar{\alpha} : K(R, n) \rightarrow X$ by dividing our arguments in two cases:

(Case 1) n is an odd positive integer > 1 , namely, $n = 2m + 1$ for some $m \geq 1$. Then we have $K(\mathbb{Q}, 2m + 1) \simeq S_{\mathbb{Q}}^{2m+1}$, and hence we have nothing to do.

(Case 2) n is an even positive integer, namely, $n = 2m$ for some $m \geq 1$. Since $\alpha_\sigma \in G(S_{\mathbb{Q}}^{2m}, X)$, the map $\alpha_\sigma : S_{\mathbb{Q}}^{2m} \rightarrow X$ can be extended to the James reduced product space by Proposition 1.2, say,

$$\bar{\alpha}_\sigma : (S_{\mathbb{Q}}^{2m})_\infty \longrightarrow X, \quad \bar{\alpha}_\sigma \in G((S_{\mathbb{Q}}^{2m})_\infty, X),$$

where we know $(S_{\mathbb{Q}}^{2m})_\infty \simeq (S_\infty^{2m})_{\mathbb{Q}} \simeq (\Omega \Sigma S^{2m})_{\mathbb{Q}} \simeq (\Omega S^{2m+1})_{\mathbb{Q}} \simeq \Omega(S_{\mathbb{Q}}^{2m+1}) \simeq \Omega K(\mathbb{Q}, 2m+1) \simeq K(\mathbb{Q}, 2m)$. Thus we have $\bar{\alpha}_\sigma \in G(K(\mathbb{Q}, 2m), X)$. Hence by Proposition 1.3, there is a map $\bar{\alpha} : K(R, 2m) = \bigoplus_{\lambda} K(\mathbb{Q}, 2m) \rightarrow X$ extending $\alpha : S^n(R) \rightarrow X$. Then we obtain $\beta \circ \bar{\alpha} \sim \text{id}_{K(R, n)}$, since the identity map $\text{id} : K(R, n) \rightarrow K(R, n)$ is the unique extension of $\iota_R^n : S^n(R) \rightarrow K(R, n)$, up to homotopy.

Thus in either case, we obtain a map $\bar{\alpha} \in G(K(R, n), X)$ such that

$$\beta \circ \bar{\alpha} \sim \text{id} : K(R, n) \longrightarrow K(R, n).$$

Let Y be the homotopy fibre of $\beta : X \rightarrow K(R, n)$. Then by Theorem 1.3 of Dula and Gottlieb [4], we obtain

$$X \simeq Y \times \bigoplus_{\lambda \in \Lambda} K(\mathbb{Q}, n) \simeq Y \times K(R, n).$$

This completes the proof of the theorem. $\square$

3. APPLICATIONS

A 0-connected space X is called a T -space if the fibration $\Omega X \rightarrow X^{S^1} \rightarrow X$ is trivial in the sense of fibre homotopy type (Aguadé [1]). If X is a 0-connected Hopf space, then X is a T -space. Aguadé showed that 1-connected space X of finite type is a rational T -space if and only if X has the same rational homotopy type as a generalized Eilenberg-Mac Lane space, i.e., a product of (infinitely many) Eilenberg-Mac Lane spaces (Theorem 3.3 of [1]). Woo and Yoon showed that a space X is a T -space if and only if $G(\Sigma A, X) = [\Sigma A, X]$ for any space A by Theorem 2.2 of [18]. So, it might be more appropriate to call such space a generalized Gottlieb space. Then we have the following result by Theorem 2.2.

Theorem 3.1. *Let $R = \bigoplus_{\lambda \in \Lambda} \mathbb{Q}$ be a finite or an infinite dimensional $\mathbb{Q}$ -vector space. Let X be a 0-connected T -space and $R \subset \pi_n(X)$, $n \geq 2$. If $\bar{p}|R : R \rightarrow H_n(X)$ is an injection, where R is represented as $R \subset [S_{\mathbb{Q}}^n, X] = G(S_{\mathbb{Q}}^n, X)$. Then X decomposes as*

$$X \simeq Y \times \bigoplus_{\lambda \in \Lambda} K(\mathbb{Q}, n) \simeq Y \times K(R, n), \quad \text{for a } T\text{-space } Y.$$

Proof. By Theorem 2.11 of [18] and Theorem 2.2, we have the result. $\square$

Theorem 3.1 implies the following result as a direct consequence.

Corollary 3.2. *Let $n \geq 2$. Let $R = \bigoplus_{\lambda \in \Lambda} \mathbb{Q}$ be a finite or an infinite dimensional $\mathbb{Q}$ -vector space and assume that $R \subset \pi_n(X)$. If X is $(n-1)$ -connected T -space, then X splits as*

$$X \simeq Y \times \bigoplus_{\lambda \in \Lambda} K(\mathbb{Q}, n) \simeq Y \times K(R, n), \quad \text{for a } T\text{-space } Y.$$

A space X is called a G -space if $G_n(X) = \pi_n(X)$ for all n (cf. [7]). As a special case of Theorem 2.2, we have the following result for rational Gottlieb space. We remark that $\pi_n(X_{\mathbb{Q}}) = G_n(X_{\mathbb{Q}})$ implies $[S_{\mathbb{Q}}^n, X_{\mathbb{Q}}] = G(S_{\mathbb{Q}}^n, X_{\mathbb{Q}})$ for any n .

Theorem 3.3. *Let $n \geq 2$. Assume that a rational space $X_{\mathbb{Q}}$ is an $(n-1)$ -connected Gottlieb space. If $\pi_n(X_{\mathbb{Q}}) \cong \bigoplus_{\lambda \in \Lambda} \mathbb{Q}$, a finite or an infinite dimensional $\mathbb{Q}$ -vector space, then $X_{\mathbb{Q}}$ decomposes as*

$$X_{\mathbb{Q}} \simeq Y_{\mathbb{Q}} \times \bigoplus_{\lambda \in \Lambda} K(\mathbb{Q}, n) \simeq Y_{\mathbb{Q}} \times K(\pi_n(X_{\mathbb{Q}}), n),$$

where $Y_{\mathbb{Q}}$ is an n -connected Gottlieb space.

Theorem 3.3 implies the following theorem (cf. [16]). For finite complexes or finite Postnikov pieces, it is known by Haslam [8] and Mataga [12].

Theorem 3.4. *If X is a 1-connected space, then the following are equivalent:*

- (1) $X_{\mathbb{Q}}$ is a Gottlieb space.
- (2) $X_{\mathbb{Q}}$ is a T -space.
- (3) $X_{\mathbb{Q}}$ is a Hopf space.
- (4) $X_{\mathbb{Q}}$ has the homotopy type of a generalized Eilenberg-Mac Lane space.

Corollary 3.5. *Any k -invariant of a 1-connected Gottlieb space is rationally trivial.*

We remark that Corollary 3.5 doesn't imply that a k -invariant of a 1-connected Gottlieb space is of finite order. Now, $H_*(K(\bigoplus_{\lambda} \mathbb{Q}, 2m+1); \mathbb{Q})$ is isomorphic to an exterior algebra and $H_*(K(\bigoplus_{\lambda} \mathbb{Q}, 2m); \mathbb{Q})$ is isomorphic to a polynomial algebra as Hopf algebras. Thus we obtain a generalization of Theorem 3.2 of Borel [2]:

Corollary 3.6. *Let X be a 1-connected rational Gottlieb space. Then $X_{\mathbb{Q}}$ is a Hopf space and the Hopf algebra $H^*(X; \mathbb{Q})$ is isomorphic (as an algebra) to the tensor product of the dual algebra of a polynomial algebra on even degree generators and the dual algebra of an exterior algebra on odd degree generators.*

We remark that $\pi_q(X) \otimes \mathbb{Q}$ may be infinite dimensional for each $q \geq 1$, and hence $H_q(X; \mathbb{Q})$ and its dual $H^q(X; \mathbb{Q}) \cong \text{Hom}(H_q(X; \mathbb{Q}); \mathbb{Q})$ may be distinct as $\mathbb{Q}$ -modules for each $q \geq 1$. For example, the dual of an exterior algebra on $\{\alpha_{\lambda}\}$ is not an exterior algebra on $\{\bar{\alpha}_{\lambda}\}$, in general, where $\bar{\alpha}_{\lambda}$ is the dual to α_{λ} (cf. [11]).

REFERENCES

- [1] J. Aguadé, *Decomposable free loop spaces*, Canad. J. Math. **39** (1987), 938–955.
- [2] A. Borel, *Topics in the homology theory of fibre bundles*, (Lecture Note in Math. **36**), Springer, Berlin 1967.
- [3] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, (Lecture notes in Math., **304**), Springer, Berlin 1972.
- [4] G. Dula and D. H. Gottlieb, *Splitting off H -spaces and Conner-Raymond splitting theorem*, J. Fac. Sci. Univ. Tokyo Sect. IA, Math. **37** (1990), 321–334.
- [5] L. Fuchs, *Infinite abelian groups*, Vol. I, (Graduate Texts in Mathematics, **61**), Academic Press, New York-London, 1970.

- [6] D. H. Gottlieb, *A certain subgroup of the fundamental group*, Amer. J. Math., **87** (1965), 840 – 856.
- [7] D. H. Gottlieb, *Evaluation subgroups of homotopy groups*, Amer. J. Math., **91** (1969), 729 – 756.
- [8] H. B. Haslam, *G-spaces mod F and H-spaces mod F*, Duke Math. J. **38**, (1971), 671–679.
- [9] P. Hilton, G. Mislin and J. Roitberg, *Localization of nilpotent groups and spaces*, (Mathematics studies, **15**), North-Holland, Amsterdam, 1975.
- [10] I. M. James, *Reduced product spaces*, Ann. of Math. **62** (1955), 170–197.
- [11] S. Mac Lane, “Homology”, Springer Verlag, Berlin, 1963.
- [12] Y. Mataga, *Localization of G-spaces*, Proc. Japan Acad. **50** (1974), 448–452.
- [13] M. Mimura, G. Nishida and H. Toda, *Localization of CW-complexes and its applications*, J. Math. Soc. Japan **23** (1971), 593–624.
- [14] N. Oda, *Pairings and copairings in the category of topological spaces*, Publ. RIMS Kyoto Univ. **28** (1992), 83–97.
- [15] J. Oprea, *Decomposition theorems in rational homotopy theory*, Proc. Amer. Math. Soc. **96** (1986), 505–512.
- [16] H. Scheerer, *On rationalized H- and co-H-spaces. With an appendix on decomposable H- and co-H-spaces*, Manuscripta Math. **51** (1985), 63–87.
- [17] K. Varadarajan, *Generalised Gottlieb groups*, J. Indian Math. Soc. **33** (1969), 141–164.
- [18] M. H. Woo and Y. S. Yoon, *T-spaces by the Gottlieb groups and duality*, J. Austral. Math. Soc. Ser. A **59** (1995), 193–203.

E-mail address: iwase@math.kyushu-u.ac.jp

E-mail address: odanobu@cis.fukuoka-u.ac.jp

(Iwase) FACULTY OF MATHEMATICS, KYUSHU UNIVERSITY, FUKUOKA 810-8560, JAPAN

(Oda) DEPARTMENT OF APPLIED MATHEMATICS, FUKUOKA UNIVERSITY, FUKUOKA 814-0180, JAPAN